



Reporting of Cancer Quality Measures: Sidney Kimmel Cancer Center - Washington Township, 2012 - 2017

To ensure continuous monitoring and reporting of quality measures, the team at the Sidney Kimmel Cancer Center - Washington Township conducts continuous practice improvements to improve the quality of patient care.

Table Key: **Green** means that our results **meet or exceed** the American College of Surgeons Commission on Cancer (COC) Benchmark - Standard 4.4/4.5. **N/A** means there were no qualifying cases.

The Commission on Cancer (CoC), a program of the American College of Surgeons (ACoS), recognizes cancer care programs for their commitment to providing comprehensive, high-quality, and multidisciplinary patient centered care.	2014	2015	2016	2017	CoC Benchmark
Tamoxifen, or third generation aromatase inhibitor, is recommended or administered within 1 year of diagnosis for women with AJCC T1c or stage IB-III hormone receptor positive breast cancer	97.10%	100%	100%	91.8%	90% Standard 4.4
Radiation is administered within 1 year of diagnosis for women under the age of 70 receiving breast conservation surgery	96.40%	100%	100%	94.7%	90% Standard 4.4
Radiation therapy is recommended or administered following any mastectomy within 1 year of diagnosis of breast cancer for women with >= 4 positive lymph nodes	N/A	100%	100%	100%	90% Standard 4.4
Image or palpation-guided needle biopsy to the primary site is performed to establish diagnosis of breast cancer	98.70%	94.10%	93.9%	100%	80% Standard 4.5
At least 12 regional lymph nodes removed and pathologically examined for resected colon cancer.	95.20%	95.70%	96.9%	100%	85% Standard 4.5

Preoperative chemo and radiation are administered for clinical AJCC T3N0, T3N0, or Stage III, or postoperative chemo and radiation are administered within 180 days of diagnosis for clinical AJCC T1-2N0 with pathological AJCC T3N0, T4N0, or Stage III; or treatment is recommended for patients under the age of 80 receiving resection for rectal cancer	100%	100%	100%	100%	85% Standard 4.5
Systemic chemotherapy is administered within 4 months to day of preoperatively, or day of surgery to 6 months postoperatively, or it is recommended for surgically resected cases with pathologic lymph node-positive (pN1) and (pN2) NSCLC	100%	100%	50% **	100%	85% Standard 4.5
Surgery is not the first course of treatment for cN2, M0 lung cases	100%	91.70%	75% **	100%	85% Standard 4.5
At least 15 regional lymph nodes removed and pathologically examined for resected gastric cancer	N/A	100%	100%	0% **	80% Standard 4.5

** This *percentage* represents a small sampling of cases that meet the specific criteria